


Arakiite



©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Tetragonal. *Point Group:* $\bar{4}2m$. As crystals composed of pyramids and prisms, to 2 mm, and in crusts.

Physical Properties: Hardness = Soft. $D(\text{meas.}) = 2.23$ $D(\text{calc.}) = 2.34$ (from cell-dimension charts). Soluble in H_2O .

Optical Properties: Translucent. *Color:* Buff to colorless. *Streak:* White. *Optical Class:* Uniaxial (-). $\omega = 1.513$ $\epsilon = 1.470$

Cell Data: *Space Group:* $I\bar{4}2d$ (synthetic KH_2PO_4). $a = 7.451(1)$ $c = 6.792(15)$ $Z = 4$

X-ray Powder Pattern: Synthetic KH_2PO_4 , (ICDD 35-807). 3.724 (100), 2.908 (83), 1.953 (51), 2.635 (23), 2.340 (15), 5.086 (14), 3.008 (14)

Chemistry:		(1)
	K_2O	10.8
	NH_3	3.46
	Total	

(1) Petrogale Cave, Western Australia; partial analysis, corresponding to $[\text{K}_{0.74}(\text{NH}_4)_{0.26}]_{\Sigma=1.00} \text{H}_2\text{PO}_4$.

Occurrence: As a component of stalactites and crusts on the walls of caves containing bat guano deposits.

Association: Biphosphammite, apthitalite, halite, syngenite, stercorite, oxammite, weddellite, whitlockite, guanine, newberyite, calcite, mundrabillaite.

Distribution: In Petrogale Cave, near Madura Motel, Eucla, and in Murra-el-elevyn Cave, Cocklebidy, Western Australia.

Name: Honors Dr. Michael Archer (1945–), Curator of Mammals, Queensland Museum, Brisbane, Australia, who discovered the first specimens.

Type Material: Western Australian Museum, Perth, Australia, MDC5901.

References: (1) Bridge, P.J. (1977) Archerite, $(\text{K}, \text{NH}_4)\text{H}_2\text{PO}_4$, a new mineral from Madura, Western Australia. *Mineral. Mag.*, 41, 33–35. (2) (1977) *Amer. Mineral.*, 62, 1057 (abs. ref. 1). (3) West, J. (1930) A quantitative X-ray analysis of the structure of potassium dihydrogen phosphate (KH_2PO_4). *Zeits. Krist.*, 74, 306–332.