


Babingtonite



©2001 Mineral Data Publishing, version 1.2

Crystal Data: Triclinic. *Point Group:* $\bar{1}$. Stubby, prismatic, striated crystals, to 3 cm; also platy and as subparallel aggregates.

Physical Properties: *Cleavage:* Perfect on {001}, good on {010} and {100}.
Fracture: Uneven. *Tenacity:* Brittle. *Hardness* = 5.5–6 *D*(meas.) = 3.34–3.37
D(calc.) = 3.26

Optical Properties: Opaque to faintly translucent. *Color:* Dark greenish black.
Luster: Vitreous.

Optical Class: Biaxial (-). *Pleochroism:* *X* = deep green; *Y* = lilac-brown; *Z* = pale to deep brown. *Dispersion:* $r > v$, strong. $\alpha = 1.700(3)$ $\beta = 1.710(3)$ $\gamma = 1.725(3)$
 $2V$ (meas.) = $50(2)^\circ$

Cell Data: *Space Group:* $P\bar{1}$. $a = 7.50(1)$ $b = 12.18(1)$ $c = 6.68(1)$ $\alpha = 86^\circ 3(5)'$
 $\beta = 93^\circ 59(5)'$ $\gamma = 112^\circ 19(5)'$ $Z = 2$

X-ray Powder Pattern: Baveno, Italy.

2.75 (100), 2.87 (80), 3.12 (70), 2.17 (60), 2.95 (60b), 2.47 (50), 6.69 (40)

Chemistry:

	(1)	(2)	(3)
SiO ₂	52.80	52.44	52.41
TiO ₂	0.29	0.00	
Al ₂ O ₃	2.41	0.23	
Fe ₂ O ₃	13.17	14.43	13.93
FeO	7.68	10.12	12.53
MnO	1.89	1.03	
MgO	0.92	0.65	
CaO	19.19	19.67	19.56
Na ₂ O	0.48	0.08	
H ₂ O	0.91	1.27	1.57
Total	99.74	[99.92]	100.00

(1) Arendal, Norway. (2) Holyoke, Hampden Co., Massachusetts, USA; original total given as 99.82%. (3) Ca₂Fe²⁺Fe³⁺Si₅O₁₄(OH).

Polymorphism & Series: Forms a series with manganbabingtonite.

Occurrence: In veins cutting granite pegmatite and diorite; in cavities and vugs in mafic volcanic rocks and gneisses; in skarns.

Association: Prehnite, calcite, epidote, albite, orthoclase, garnet, quartz, "hornblende," zeolites.

Distribution: Known from a number of localities, most minor; some providing good crystals are: in Norway, at the Brastad mine, Oyestad, near Arendal. From Gronsjöberg, Dalarma, Sweden. In Germany, at Herbornseelbach, Hesse. From Baveno, Piedmont, Italy. In the Khandivali quarry, near Bombay, Maharashtra, India. At Mitani, Kochi Prefecture, Japan. From Noril'sk, western Siberia, Russia. In the USA, in Massachusetts, exceptional crystals from Lane's quarry, Westfield, Hampden Co.; in the Cheapside and Deerfield quarries, East Deerfield, Franklin Co.; also at Winchester Highlands, Uxbridge, Norfolk Co. At Paterson, Passaic Co., and Mine Hill, Morris Co., New Jersey; from the Goose Creek quarry, Leesburg, Loudoun Co., Virginia; and at Lenoir, Caldwell Co., North Carolina.

Name: For Dr. William Babington (1756–1833), Irish physician and mineralogist.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.

References: (1) Dana, E.S. (1892) Dana's system of mineralogy, (6th edition), 381–382. (2) Washington, H.S. and H.E. Merwin (1923) On babingtonite. *Amer. Mineral.*, 8, 215–223. (3) Palache, C. and F.A. Gonyer (1932) On babingtonite. *Amer. Mineral.*, 17, 295–299. (4) Kosoi, A.L. (1975) The structure of babingtonite. *Kristallografiya (Sov. Phys. Crystal.)*, 20, 730–739 (in Russian). (5) Traill, R.J. and A.P. Sabina (1960) Catalogue of X-ray diffraction patterns and specimen mounts on file at the Geological Survey of Canada. *Geol. Survey of Canada, Paper 60-4*, 11.