Goslarite $ZnSO_4 \cdot 7H_2O$

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic. *Point Group:* 222. Typically stalactitic or in efflorescent crusts; fibrous, granular or massive.

Physical Properties: Cleavage: On $\{010\}$, perfect. Tenacity: Brittle. Hardness = 2–2.5 D(meas.) = 1.98(5) D(calc.) = 1.972 Soluble in H_2O , taste astringent, metallic, nauseous; dehydrates in air; strongly diamagnetic.

Optical Properties: Transparent. *Color:* Colorless, white; pale shades of blue, green, or brown from impurities; colorless in transmitted light. *Luster:* Vitreous, silky if fibrous. *Optical Class:* Biaxial (–). *Orientation:* X = b; Y = c; Z = a. *Dispersion:* r > v, weak. $\alpha = 1.463$ $\beta = 1.475$ $\gamma = 1.480$ 2V(meas.) = Small to moderate.

Cell Data: Space Group: $P2_12_12_1$. a = 11.779 b = 12.050 c = 6.822 Z = 4

X-ray Powder Pattern: Synthetic.

4.21 (100), 5.36 (80), 4.18 (50), 5.29 (35), 2.867 (35), 2.651 (30), 4.46 (25)

	(1)	(2)
SO_3	27.29	27.84
FeO	1.12	
MnO	1.24	
NiO	0.06	
ZnO	23.12	28.30
MgO	0.81	
H_2O	42.42	43.86
insol.	3.65	
Total	99.71	100.00

(1) Ushaw Moor, Durham, England. (2) $ZnSO_4 \cdot 7H_2O$.

Polymorphism & Series: Forms a series with epsomite.

Occurrence: A secondary mineral in oxidized portions of zinc sulfide deposits, particularly as coatings on walls of mine passages.

Association: Epsomite, melanterite, pickeringite, chalcanthite, gypsum.

Distribution: Widespread as post-mining efflorescences. A few notable localities include: in Germany, from the Rammelsberg mine, near Goslar, Harz Mountains, and at Freiberg and Altenberg, Saxony. From Baňská Štiavnica (Schemnitz), Slovakia. At Falun, Sweden. From Sain-Bel, near Lyon, Rhône, France. At the Almagrera mine, Tharsis, Huelva Province, Spain. In Italy, at Capanne Vecchie, Elba, and Raibl, Venezia Giulia. At Broken Hill, New South Wales, Australia. In the USA, in Arizona, at Bisbee, Cochise Co. and many other districts; from the Central district, Grant Co., New Mexico; at Island Mountain, Trinity Co., California; from Butte, Silver Bow Co., Montana; at Bingham Canyon, Salt Lake Co., Utah; in the Comstock Lode, Storey Co., Nevada; at Friedensville, Lehigh Co., Pennsylvania. From Quebrada de La Alcaparrosa, San Juan Province, Argentina.

Name: For the locality at Goslar, Germany.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 513–516. (2) (1959) NBS Circ. 539, 8, 71–73.