Crystal Data: Triclinic. Point Group: 1 or $\overline{1}$. As fibrous aggregates, may be plumose, to 4.5 cm; individuals are elongated along [001]. Twinning: An indeterminate twinninglike intergrowth was observed.

Cleavage: Good on {100}, poor on {010}. Tenacity: Friable in Physical Properties: aggregate. Hardness = n.d. D(meas.) = 3.48(4) D(calc.) = 3.51

Optical Properties: Transparent. Color: Light yellowish brown; colorless in transmitted light. Streak: Light brownish yellow. Luster: Vitreous.

Optical Class: Biaxial (-). Dispersion: r > v, distinct. $\alpha = 1.6742(4)$ $\beta = 1.6968(3)$ $\gamma = 1.6999(3)$ 2V(meas.) = 41.9(2)° 2V(calc.) = 40.9°

Cell Data: Space Group: P1 or $P\overline{1}$. a = 10.44(2) b = 11.064(6) c = 9.62(1) $\alpha = 107.43(7)^{\circ}$ $\beta = 82.7(1)^{\circ}$ $\gamma = 111.6(1)^{\circ}$ Z = 1

X-ray Powder Pattern: Kombat mine, Namibia. 2.676 (100), 9.8 (60), 3.23 (60), 5.99 (40), 3.38 (40), 2.479 (40), 1.539 (40b)

Chemistry:

	(1)	(2)
SiO_2	35.5	34.68
$\mathrm{As_2O_5}$	10.6	11.05
FeO	0.1	
MnO	40.7	40.94
$_{\rm MgO}$	8.2	7.75
Na_2O	3.1	2.98
$\mathrm{H_2O}$	2.6	2.60
Total	100.8	100.00

(1) Kombat mine, Namibia; by electron microprobe, total As as As₂O₅, total Mn as MnO, H₂O by the Penfield method. (2) $Na_2Mg_4Mn_{12}As_2Si_{12}O_{43}(OH)_6$.

Occurrence: In low-temperature hydrothermal veins cutting metamorphosed Fe-Mn ores.

Association: Kentrolite, rhodonite, richterite, barite, calcite.

Distribution: From the Kombat mine, 49 km south of Tsumeb, Namibia.

Name: Honors John Innes, mineralogist of the Tsumeb Corporation, for his contributions to the mineralogy of the Tsumeb and Kombat mines.

Type Material: National Museum of Natural History, Washington, D.C., USA, 163207.

References: (1) Dunn, P.J., D.R. Peacor, S.-C. Su, J.A. Nelen, and O. von Knorring (1986) Johninnesite, a new sodium manganese arsenosilicate from the Kombat Mine, Namibia. Mineral. Mag., 50, 667-670. (2) (1988) Amer. Mineral., 73, 928 (abs. ref. 1).