

**Crystal Data:** Monoclinic, pseudorhombic. *Point Group:* 2/m. Crystals are platy or bladed prismatic, to 1 mm; typically in dendritic, reticulated, stellate, or fibrous aggregates; also fine granular massive. *Twinning:* Reticulated groups, not described as twins, are characteristic.

**Physical Properties:** *Fracture:* Uneven. *Tenacity:* Brittle. Hardness = ~4.5 VHN = 250–320, 280 average (25 g load). D(meas.) = 4.71 D(calc.) = 4.62–4.73

**Optical Properties:** Transparent to translucent, may be opaque from inclusions.

*Color:* Colorless, white, silvery gray, pale pink, yellow, greenish yellow, reddish brown.

*Streak:* White. *Luster:* Vitreous to greasy.

*Optical Class:* Biaxial (-). *Orientation:* Y = b; Z ∧ c = 26.5°. *Dispersion:* r > v, medium. α = 1.584–1.594 β = 1.710–1.730 γ = 1.715–1.730 2V(meas.) = 10°–20° 2V(calc.) = 18°–22°

**Cell Data:** *Space Group:* P2<sub>1</sub>/m. a = 13.360–13.418 b = 5.052–5.077 c = 6.681–6.714 β = 106.56°–106.58° Z = 2

**X-ray Powder Pattern:** Khibiny massif, Russia.

4.00 (100), 3.269 (100), 2.140 (40), 2.003 (40), 2.535 (20), 1.635 (10), 1.373 (10)

Chemistry:	(1)	(2)	(1)	(2)
CO <sub>2</sub>	[21.95]	21.10	CaO	0.39
Y <sub>2</sub> O <sub>3</sub>	0.15		SrO	1.55
(La, Nd) <sub>2</sub> O <sub>3</sub>		13.89	BaO	47.39
La <sub>2</sub> O <sub>3</sub>	6.61		F	3.18
Ce <sub>2</sub> O <sub>3</sub>	15.30	10.96	–O = F <sub>2</sub>	1.34
Pr <sub>2</sub> O <sub>3</sub>	1.19		rem.	2.99
Nd <sub>2</sub> O <sub>3</sub>	4.26		Total	[100.63]
MgO		0.85		100.03

(1) Khibiny massif, Russia; by electron microprobe, average of 12 analyses, F by microchemical methods, CO<sub>3</sub><sup>2-</sup> calculated, corresponding to (Ba<sub>1.86</sub>Sr<sub>0.09</sub>Ca<sub>0.04</sub>)<sub>Σ=1.99</sub>(Ce<sub>0.56</sub>La<sub>0.24</sub>Nd<sub>0.15</sub>Pr<sub>0.04</sub>Y<sub>0.01</sub>)<sub>Σ=1.00</sub>(CO<sub>3</sub>)<sub>3.00</sub>F<sub>1.01</sub>. (2) Bayan Obo deposit, China; remainder is ThO<sub>2</sub> 0.41%, Al<sub>2</sub>O<sub>3</sub> 0.06%, Fe<sub>2</sub>O<sub>3</sub> 2.52%; corresponds to (Ba<sub>1.56</sub>Ca<sub>0.22</sub>Mg<sub>0.11</sub>Sr<sub>0.09</sub>)<sub>Σ=1.98</sub>(Ce<sub>0.34</sub>Ln<sub>0.46</sub>)<sub>Σ=0.80</sub>(CO<sub>3</sub>)<sub>3</sub>F.

**Occurrence:** In syenitic alkalic massifs and carbonatites.

**Association:** Belovite-(Ce), ancylite-(Ce), cebaite-(Ce), cordylite-(Ce), synchysite-(Ce), mckelveyite-(Y), ewaldite, rinkite, zakharovite, nenadkevichite, yofortierite, lävenite, burbankite, calcite, barite, fluorapatite, fluorite.

**Distribution:** In Russia, from Tuliylukht Bay and the Kirov apatite mine, Mt. Kukisvumchorr, Khibiny massif, and at the Vuoriyarvi carbonatite complex, Kola Peninsula. In the Bayan Obo Fe–Nb–Re deposit, 130 km north of Baotou, Inner Mongolia, China. At Mont Saint-Hilaire, and from near Saint-Amable, Quebec, Canada.

**Name:** To honor Professor Aleksandr Aleksandrovich Kukharenko (1914–1993), Russian mineralogist, St. Petersburg University, St. Petersburg, Russia, for his extensive work on the carbonatites of the Kola Peninsula, Russia.

**Type Material:** Geology Museum, Kola Branch, Academy of Sciences, Apatity, 6097; Mineralogical Museum, St. Petersburg University, Russia, 1/18303; Canadian Museum of Nature, Ottawa, Canada, 81531–81533.

**References:** (1) Zaitsev, A.N., V.N. Yakovenchuk, G.Y. Chao, R.A. Gault, V.V. Subbotin, Y.A. Pakhomovskii, and A.N. Bogdanova (1996) Kukharenkoite-(Ce), Ba<sub>2</sub>Ce(CO<sub>3</sub>)<sub>3</sub>F, a new mineral from Kola peninsula, Russia, and Québec, Canada. *Eur. J. Mineral.*, 8, 1327–1336. (2) (1997) *Amer. Mineral.*, 82, 1039 (abs. ref. 1). (3) Zhang Peishan and Tao Kejie (1981) Zhonghuacerite [= kukharenkoite-(Ce)] Ba<sub>2</sub>Ce(CO<sub>3</sub>)<sub>3</sub>F – a new mineral. *Scientia Geologica Sinica*, 195–196 (in Chinese with English abs.). (4) Mercier, N. and M. Leblanc (1993) Crystal growth and structures of rare earth fluorocarbonates: II. Structure of zhonghuacerite [= kukharenkoite-(Ce)] Ba<sub>2</sub>Ce(CO<sub>3</sub>)<sub>3</sub>F. *Eur. J. Solid State Inorg. Chem.*, 30, 207–216.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.