(c)2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic, pseudohexagonal. *Point Group:* 222 (probable). Acicular to elongated lozenge-shaped crystals, to 5 μ m; may form star-shaped aggregates.

Physical Properties: Tenacity: Friable. Hardness = n.d. D(meas.) = > 3.30 D(calc.) = [4.70] Radioactive.

Optical Properties: Semitransparent. Color: Brownish green to brown. Optical Class: Biaxial; low birefringence. Pleochroism: Slight; X = pale brown. Orientation: Parallel extinction; positive elongation. n = 1.60-1.70

Cell Data: Space Group: P222 (probable). a = 6.78(3) b = 12.10(5) c = 6.38(3) Z = 3

X-ray Powder Pattern: Ningyo-Toge mine, Japan. 3.02 (vsb), 2.81 (s), 2.13 (s), 4.33 (m), 3.38 (mb), 5.99 (wb), 3.45 (w)

Chemistry:

	(1)
UO_3	n.d.
P_2O_5	16.8
$\overline{\mathrm{UO}_{2}}$	23.3
FeO	4.8
CaO	6.1
$\mathrm{H_2O^+}$	7.4
$\overline{\mathrm{H}_{2}\mathrm{O}^{-}}$	1.9
$\overline{\mathrm{H}_{2}\mathrm{O}}$	
\overline{C}	2.3
insol.	30.9
Total	93.5

(1) Ningyo-Toge mine, Japan; RE \approx a few %, which may be in included a patite, corresponds to $\mathrm{Ca_{0.92}U_{0.73}Fe_{0.56}(PO_4)_2} \bullet 3.47\mathrm{H_2O}$.

Mineral Group: Rhabdophane group.

Occurrence: The principal uranium mineral, as coatings and filling cavities in carbonaceous conglomerate (Ningyo mine, Japan); in uranium deposits in organic-rich conglomerate-sandstone-mudstone (British Columbia, Canada).

Association: Pyrite, marcasite, sphalerite, apatite, chlorite, gypsum, quartz, hypersthene, feldspar, clay minerals (Ningyo mine, Japan); saléeite, autunite, uraninite (Okanagan district, Canada).

Distribution: From the Ningyo-Toge uranium mine, Tottori Prefecture, Japan. In the Tyee and Blizzard uranium deposits, 60 and 30 km southeast of Kelowna, Okanagan district, British Columbia, Canada. From the Dzhantuar and Rudnoye uranium deposits, Auminzatau Mountains, central Kyzylkum region, Uzbekistan. At an undefined locality in the Czech Republic.

Name: For Ningyo Pass, Japan, where the mine in which the mineral was first found is located.

Type Material: The Natural History Museum, London, England, 1960,92; National Museum of Natural History, Washington, D.C., USA, 113822, 115885.

References: (1) Muto, T., R. Meyrowitz, A.M. Pommer, and T. Murano (1959) Ningyoite, a new uranous phosphate mineral from Japan. Amer. Mineral., 44, 633–650. (2) Boyle, D.R., A.L. Littlejohn, A.C. Roberts, and D.M. Watson (1981) Ningyoite in uranium deposits of south-central British Columbia: first North American occurrence. Can. Mineral., 19, 325–331.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.