(c)2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** Tetragonal. Point Group:  $4/m \ 2/m \ 2/m$ ,  $\overline{4}2m$ , 4mm, or 422. As bladed crystals, to 0.3 mm, and crystalline aggregates, embedded in other lead minerals.

**Physical Properties:** Cleavage:  $\{001\}$ , perfect;  $\{100\}$ , possible, fair. Tenacity: Sectile. Hardness = 2–2.5 VHN = 113–133, 122 average (50 g load). D(meas.) = 7.32 (synthetic). D(calc.) = 7.39

**Optical Properties:** Translucent. *Color:* Bright red, purplish red, brownish red; brownish red in transmitted light; in reflected light, gray with deep red to red-orange internal reflections. *Streak:* Scarlet to grenadine-red. *Luster:* Adamantine, pearly to resinous.

Optical Class: Uniaxial (–); anomalously biaxial. Pleochroism: Strong; O = brownish red; E = greenish yellow.  $\omega = [2.58]$   $\epsilon = [2.42]$  2V(meas.) = Small. Anisotropism: Weak to moderate. Bireflectance: Moderate.

 $\begin{array}{l} R_1-R_2\colon (400)\ 22.6-21.8, (420)\ 22.4-21.2, (440)\ 22.3-20.5, (460)\ 22.2-19.9, (480)\ 21.9-19.4, (500)\ 21.6-18.9, (520)\ 21.2-18.5, (540)\ 20.6-18.1, (560)\ 20.2-17.8, (580)\ 19.8-17.5, (600)\ 19.4-17.2, (620)\ 19.1-17.1, (640)\ 18.9-17.0, (660)\ 18.7-16.9, (680)\ 18.6-16.7, (700)\ 18.5-16.7 \end{array}$ 

**Cell Data:** Space Group: I4/mmm,  $I\overline{4}2m$ ,  $I\overline{4}m2$ , I4mm, or I422. a = 3.9922(3) c = 22.514(3) Z = 2

X-ray Powder Pattern: Merehead quarry, England. 2.983 (100), 2.816 (78), 1.989 (75), 1.658 (51), 1.586 (33), 3.507 (32), 1.263 (24)

Chemistry:

$$\begin{array}{ccc} & & & (1) \\ \text{MoO}_3 & 8.0 \\ \text{PbO} & 88.4 \\ \text{Cl} & 4.4 \\ -\text{O} = \text{Cl}_2 & 1.0 \\ \hline \text{Total} & [99.8] \end{array}$$

(1) Merehead quarry, England; by electron microprobe, average of six analyses, original total given as 100.8%; corresponds to  $(Pb_{3.17}Mo_{0.44})_{\Sigma=3.61}O_{4.01}Cl_{0.99}$ .

**Occurrence:** A very rare mineral in nodules in manganese oxide ores in veins, fissures, and pockets in limestone.

**Association:** Mendipite, chloroxiphite, diaboleite, cerussite, paralaurionite, wulfenite, calcite, manganese oxides (Merehead quarry, England).

**Distribution:** In the Merehead quarry, Cranmore, and from the Wesley mine, Didsbury, near Bristol, Somerset, England.

Name: For Reginald F.D. Parkinson (1928–), British mineral collector and dealer, Shepton Mallet, Somerset, England, who first found the mineral.

**Type Material:** The Natural History Museum, London, England, 1987,478–479.

**References:** (1) Symes, R.F., G. Cressey, A.J. Criddle, C.J. Stanley, J.G. Francis, and G.C. Jones (1994) Parkinsonite, (Pb, Mo,  $\square$ )<sub>8</sub>O<sub>8</sub>Cl<sub>2</sub>, a new mineral from Merehead Quarry, Somerset. Mineral. Mag., 58, 59–68. (2) (1994) Amer. Mineral., 79, 1011 (abs. ref. 1).