

Raite

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Orthorhombic. *Point Group:* 222. Crystals acicular, to 2 mm; in radiating clusters.

Physical Properties: Cleavage: Perfect on {100}, {010}, and {001}. Tenacity: Brittle. Hardness = 3 D(meas.) = 2.32–2.39 D(calc.) = [2.51]

Optical Properties: Semitransparent. Color: Gold, brown, reddish brown, yellow, tan, rose, lavender, bronze. Streak: Yellow. Luster: Vitreous, may be silky.

Optical Class: Biaxial (+). Pleochroism: Distinct; X = nearly colorless; Y = yellowish; Z = golden brown. Orientation: $Z \wedge c = 0^\circ\text{--}30^\circ$. $\alpha = 1.540(2)$ $\beta = 1.542(2)$ $\gamma = 1.550(2)$ 2V(meas.) = n.d. 2V(calc.) = 53°

Cell Data: Space Group: C222. $a = 30.44\text{--}30.6$ $b = 5.31\text{--}5.371$ $c = 18.20\text{--}18.25$ $Z = 4$

X-ray Powder Pattern: Lovozerovo massif, Russia.
11.4 (100), 2.939 (100), 2.650 (100), 4.5 (80), 3.8 (60b), 2.482 (60), 1.640 (60)

Chemistry:

	(1)		(1)
SiO ₂	46.20	MnO	15.00
TiO ₂	3.11	MgO	0.20
ZrO ₂	0.16	CaO	1.24
Al ₂ O ₃	0.12	Na ₂ O	11.24
RE ₂ O ₃	0.16	K ₂ O	0.17
Fe ₂ O ₃	1.86	H ₂ O ⁺	8.01
(Nb, Ta) ₂ O ₅	0.44	H ₂ O ⁻	11.36
FeO	0.37	CO ₂	0.40
Total	100.04		

(1) Lovozerovo massif, Russia.

Occurrence: On the walls of fractures filled with nepheline in alkalic pegmatite in a differentiated alkalic massif (Lovozerovo massif, Russia).

Association: Nepheline, aegirine, mountaintite, natrolite, zorite (Lovozerovo massif, Russia); aegirine, albite, nepheline, sodalite, sérandite, analcime, ancyllite, epididymite, eudialyte, nenankevichite (Mont Saint-Hilaire, Canada).

Distribution: From the Jubilee pegmatite, on Mt. Karnasurt, near the valley of the Ilmajok River, Lovozerovo massif, and in the Khibiny massif, Kola Peninsula, Russia. At Mont Saint-Hilaire, Quebec, Canada.

Name: In honor of the international group of scientists, captained by Thor Heyerdahl, crewing the papyrus ship *Ra* (1969–1970).

Type Material: Geology Museum, Kola Branch, Academy of Sciences, Apatity, 3206, 3271; Mineralogical Museum, St. Petersburg University, St. Petersburg, 19047; Mining Institute, St. Petersburg, 1060/1–4; A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia, 74489.

References: (1) Mer'kov, A.N., I.V. Bussen, E.A. Goiko, E.A. Kul'chitskaya, Y.P. Men'shikov, and A.P. Nedorezova (1973) Raite and zorite, new minerals from the Lovozerovo Tundra. *Zap. Vses. Mineral. Obshch.*, 102, 54–62 (in Russian). (2) (1973) Amer. Mineral., 58, 1113–1114 (abs. ref. 1). (3) Khomyakov, A.P., E.M. Es'kova, G.E. Cherepivskaya, V.V. Kaptsov, and A.D. Timchenko (1982) New data on raite. *Nov. Dannye Miner.*, 30, 205–207 (in Russian). (4) (1983) Chem. Abs., 219080 (abs. ref. 3). (5) Mandarino, J.A. and V. Anderson (1989) Monteregian treasures. Cambridge Univ. Press, 174.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.