©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic (probable); metamict. *Point Group:* n.d. Crystals rough, showing three pinacoids, a prism, and an orthodome.

Physical Properties: Fracture: Uneven, small conchoidal. Hardness = 5.5-6 D(meas.) = 4.80 D(calc.) = n.d. Radioactive.

Optical Properties: Opaque, transparent in thin fragments. Color: Black, brown where altered; in transmitted light, amber-yellow to light yellow. Luster: Bright pitchy to resinous. Optical Class: Isotropic; weakly anisotropic in part. n = 2.12-2.15

Cell Data: Space Group: n.d. Z = n.d.

X-ray Powder Pattern: n.d.

Chemistry:		(1)	
	UO_3	0.64	$(\mathrm{Ce},\mathrm{La})_2\mathrm{O}_3$

UO_3	0.64	$(Ce, La)_2O_3$	2.92
$\widetilde{WO_3}$	1.87	$\mathrm{Fe_2O_3}$	1.44
Nb_2O_5	trace	$\overline{\mathrm{MnO}}$	0.13
Ta_2O_5	trace	PbO	0.48
SiO_2	trace	$_{ m MgO}$	trace
TiO_2	49.72	CaO	1.83
$\overline{\text{ThO}_2}$	8.75	$\mathrm{H_{2}O^{+}}$	4.36
UO_2	1.98	$H_2^-O^-$	0.10
$(Y, Er)_2O_3$	25.67	$\tilde{\text{CO}}_2$	0.68

 $\frac{\text{CO}_2}{\text{Total}}$ $\frac{0.68}{100.57}$

(1)

(1) Burnet Co., Texas, USA.

Occurrence: In granite pegmatite.

Association: n.d.

Distribution: In the USA, from Burnet Co., Texas, about five km east of Baringer Hill.

Name: From a dominance of YTTRium in the composition, and the Greek for a mixing, as there are many other elements in addition to yttrium.

Type Material: National Museum of Natural History, Washington, D.C., USA, 119841.

References: (1) Palache, C., H. Berman, and C. Frondel (1944) Dana's system of mineralogy, (7th edition), v. I, 793.